

The Status of V_{ud}

J.C. Hardy*

Cyclotron Institute, Texas A&M University, College Station, TX 77843

The best value for V_{ud} comes from superallowed $0^+ \rightarrow 0^+$ nuclear β transitions, of which thirteen have now been measured with high precision. The current status of these measurements is described, and the result compared with that of measurements from neutron decay and pion β decay. Future prospects for improvement are discussed.

I. INTRODUCTION

The up-down quark mixing element, V_{ud} , of the Cabibbo Kobayashi Maskawa (CKM) matrix can be determined via three β -decay routes: superallowed $0^+ \rightarrow 0^+$ nuclear decays, neutron decay and pion decay. Historically, the nuclear decays have yielded the most precise measurements, but the resultant value for V_{ud} was often thought to be severely limited in accuracy by the calculated nuclear-structure-dependent corrections that were required to extract it from the data. The neutron measurements, though free of nuclear corrections, are experimentally challenging and have therefore been less precise, not to mention occasionally inconsistent with one another. Pion β decay, being a 10^{-8} decay branch, is even more challenging experimentally and has produced larger uncertainties for V_{ud} than either of the other two approaches.

All three decay modes require that small radiative corrections be applied to the primary experimental data in the process of obtaining a value for V_{ud} , and naturally there are uncertainties associated with these calculated corrections. In fact, in the case of the superallowed nuclear decays, the experimental uncertainties have become so well controlled that it is these theoretical uncertainties that dominate the uncertainty quoted on V_{ud} . However, it may be a surprise to some readers that the nuclear-structure-dependent correction is *not* the main contributor. Instead, the dominant theoretical uncertainty originates from the so-called inner radiative correction, which is a correction that is common to all three decay modes and, unless it is improved, will ultimately limit the precision with which V_{ud} can be determined by any route.

As more and more superallowed $0^+ \rightarrow 0^+$ nuclear transitions are measured with high precision, the nuclear-structure-dependent corrections continue to prove their validity. The calculations themselves are based on well-established nuclear structure information derived from nuclear measurements that are totally independent of the superallowed decay experiments. The magnitude of the calculated correction for each transition, though always less than 1.5%, differs considerably from transition to transition. The measured superallowed transition strengths, which are an order of magnitude more pre-

cise than that, actually reproduce these predicted differences and, as a result, lead to completely consistent values of V_{ud} . All evidence points to the nuclear-structure-dependent corrections being completely reliable within their quoted uncertainties.

II. SUPERALLOWED NUCLEAR BETA DECAY

Beta decay between nuclear analog states of spin-parity, $J^\pi = 0^+$, and isospin, $T = 1$, is a pure vector transition and is nearly independent of the nuclear structure of the parent and daughter states. The measured strength of such a transition – expressed as an “ ft value” – can then be related directly to the vector coupling constant, G_V with the intervention of only a few small ($\sim 1\%$) calculated terms to account for radiative and nuclear-structure-dependent effects. Once G_V has been determined in this way, it is only another short step to obtain a value for V_{ud} , the up-down mixing element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix.

In dealing with these decays, which are referred to as superallowed, it is convenient to combine some of the small correction terms with the measured ft -value and define a “corrected” $\mathcal{F}t$ -value. Thus, we write [1]

$$\mathcal{F}t \equiv ft(1 + \delta'_R)(1 + \delta_{NS} - \delta_C) = \frac{K}{2G_V^2(1 + \Delta_R^V)}, \quad (1)$$

where $K/(\hbar c)^6 = 2\pi^3 \hbar \ln 2 / (m_e c^2)^5 = 8120.271(12) \times 10^{-10} \text{ GeV}^{-4}\text{s}$; δ_C is the isospin-symmetry-breaking correction and Δ_R^V is the transition-independent part of the radiative correction – the “inner” radiative correction. The terms δ'_R and δ_{NS} comprise the transition-dependent part of the radiative correction, the former being a function only of the electron’s energy and the Z of the daughter nucleus, while the latter, like δ_C , depends in its evaluation on the details of nuclear structure. From this equation, it can be seen that a measurement of any one of these superallowed transitions establishes an individual value for G_V ; moreover, if the Conserved Vector Current (CVC) assertion is correct that G_V is not renormalized in the nuclear medium, all such values – and all the $\mathcal{F}t$ -values themselves – should be identical within uncertainties, regardless of the specific nuclei involved.

This assertion of CVC can be tested and a value for G_V obtained with a precision considerably better than 0.1% if experiment can meet the challenge, since the four

*Electronic address: hardy@comp.tamu.edu




FIG. 1: Results from the 2005 survey [1] updated with more recent published results [2, 3, 4, 5, 6, 7, 8, 9]. The uncorrected ft values for the thirteen best known superallowed decays (left) are compared with the same results after corrections have been applied (right). The grey band in the right-hand panel is the average $\mathcal{F}t$ value, including its uncertainty.

small corrections terms only contribute to the overall uncertainty at the 0.03% level. As it turns out, experiment has exceeded that goal, leaving theory currently as the dominant contributor to the uncertainty.

The ft -value that characterizes any β -transition depends on three measured quantities: the total transition energy, Q_{EC} ; the half-life, $t_{1/2}$, of the parent state; and the branching ratio, R , for the particular transition of interest. The Q_{EC} -value is required to determine the statistical rate function, f , while the half-life and branching ratio combine to yield the partial half-life, t . In 2005 a new survey of world data on superallowed $0^+ \rightarrow 0^+$ beta decays was published [1]. All previously published measurements were included, even those that were based on outdated calibrations if enough information was provided that they could be corrected to modern standards. In all, more than 125 independent measurements of comparable precision, spanning four decades, made the cut. Another 8 publications, with data that can also be incorporated, have appeared [2, 3, 4, 5, 6, 7, 8, 9] in the two years since the survey was closed and we have now obtained ft and corrected $\mathcal{F}t$ values for the combined data¹. These updated results for the thirteen most precisely known transitions are shown in Fig. 1.

By comparing the left and right panels in Fig. 1, we can see that the calculated corrections act very well to remove

the considerable scatter that is evident in the former but missing in the latter. Since δ'_R has essentially the same value for all transitions other than those from ^{10}C and ^{14}O , the removal of the scatter must be attributed to the calculated nuclear-structure-dependent corrections, δ_{NS} and δ_C , that have been applied in the derivation of $\mathcal{F}t$. The calculation of these correction terms [10] was based on nuclear-structure models that were solidly grounded in a wide variety of independent nuclear measurements, none involving superallowed β decay. Thus the consistency of the $\mathcal{F}t$ values is a powerful validation of those calculated corrections. All thirteen $\mathcal{F}t$ values form a statistically consistent set – with a normalized chi-square of 0.9 – over a wide range of nuclear masses, from $A=10$ to $A=74$, a conclusion that is entirely consistent with CVC expectation. The average of all thirteen cases is $\overline{\mathcal{F}t} = 3073.9(8)\text{s}$. Since $\mathcal{F}t$ is inversely proportional to the square of G_V , this result confirms the constancy of the latter to 1.3 parts in 10^4 , the tightest limit ever set.

The 2005 survey results were also used to set a limit on any possible contribution from scalar currents. The presence of a scalar current – induced or fundamental – would manifest itself as a curvature, either upwards or downwards, in the $\mathcal{F}t$ -value line at low Z . There is no hint of any such curvature in the right panel of Fig.1, and a careful analysis of the survey results set a limit [1] on the scalar relative to the vector current of $|C_S/C_V| \leq 0.0013$, again the tightest limit ever set.

With a mutually consistent set of $\mathcal{F}t$ values, no scalar currents, and the test of CVC passed, one can confidently proceed to determining the value of G_V and, from it, the up-down element of the CKM matrix via the relation $V_{ud} = G_V/G_F$, where G_F is the well known [12] weak-interaction constant for purely leptonic muon decay. From the $\mathcal{F}t$ -value data in Fig. 1 and the recently

¹ In calculating the $\mathcal{F}t$ values, we have used slightly different values for δ'_R and δ_{NS} than those used by us previously [10]. This is because the recent improvements in Δ_R^V by Marciano and Sirlin [11] isolated some transition-dependent components which were more appropriately included in these other terms. Although the average value of $\mathcal{F}t$ is thus changed by $\sim 0.3\%$, a corresponding change in Δ_R^V leaves the central value of V_{ud} essentially unchanged.




FIG. 2: Error budgets for the three different methods to determine V_{ud} , illustrating the relative importance of experimental uncertainties and theoretical ones. The unbracketed values of V_{ud} correspond to the results shown in numbered equations in the text; bracketed values are alternative results obtained under different conditions also described in the text.

improved calculation of Δ_R^V [11] we obtain the result,

$$V_{ud} = 0.97378(27). \quad [\text{nuclear}] \quad (2)$$

This number can be compared with our previous value, quoted in the 2005 survey, $V_{ud} = 0.97380(40)$. The new result is completely consistent with the earlier one, but it has a considerably reduced uncertainty – mostly due to the improvement in the calculated “inner” radiative correction, Δ_R^V .

The “error budget” leading to the uncertainty quoted for V_{ud} in Eq. 2 is illustrated in the left panel of Figure 2. The largest contribution to the uncertainty comes from Δ_R^V (± 0.00018), while the nuclear-structure-dependent corrections contribute ± 0.00015 and experimental uncertainties only ± 0.00007 .

III. NEUTRON AND PION BETA DECAYS

On the one hand, free neutron decay has an advantage over nuclear decays since there are no nuclear-structure-dependent corrections to be calculated. On the other hand, it has the disadvantage that it is not purely vector-like but has a mix of vector and axial-vector contributions. Thus, in addition to a lifetime measurement, a correlation experiment is also required to separate the vector and axial-vector pieces. Both types of experiment present serious experimental challenges. The value of V_{ud} is determined from the expression

$$V_{ud}^2 = \frac{K/\ln 2}{G_F^2(1 + \Delta_R^V)(1 + 3\lambda^2)f(1 + \delta_R)\tau_n}, \quad (3)$$

where λ is the ratio of axial-vector and vector effective coupling constants, τ_n is the mean life for neutron decay,

f is the statistical rate function and δ_R is the transition-dependent radiative correction evaluated for the case of a neutron. Both λ and τ_n must be determined from experiment.

In 2003, a survey of world data for neutron β decay was published [13], in which the same criteria were used as for the superallowed nuclear beta-decay measurements: *i.e.* all measurements were retained unless they had been withdrawn by the original authors or could be rejected on objective grounds. Since that time, there have been two new mean-life measurements [14, 15] that fit within the criteria for inclusion. Strikingly, one of these measurements, by Serebrov *et al.* [15], disagrees with the previous world average by more than six standard deviations. However, without any objective grounds for rejecting this discrepant result – or, alternatively, rejecting all preceding measurements – we have evaluated the world average with all data included. The result for the neutron mean life is then $\tau_n = 882.0(15)$ s (with $\chi^2/N = 5.4$); and for the coupling-constant ratio is $\lambda = -1.2690(28)$ (with $\chi^2/N = 2.6$).

Inserting these average experimental results for τ_n and λ into Eq. 3, we determine V_{ud} to be

$$V_{ud} = 0.9766(20), \quad [\text{neutron}] \quad (4)$$

a value with much larger uncertainty than the nuclear result and just outside of one standard deviation away from it. It is also worth noting, though, that in considering the neutron mean life, the Particle Data Group has taken a different approach [12] and has completely rejected the Serebrov *et al.* [15] result simply because it does disagree significantly with previous measurements. If we do the same thing, then we obtain $V_{ud} = 0.9747(18)$, which fully overlaps the nuclear result but still has an uncertainty that is nearly 7 times larger.

The error budget for the result in Eq. 4 appears in the center panel of Figure 2. Unlike the nuclear result in the left panel, this uncertainty is completely dominated by experiment. Nevertheless, it is important to recognize that the contribution from Δ_R^V is the same for the neutron as it is for the nuclear decays. Even as the neutron’s experimental results are improved, the value of V_{ud} obtained from them will ultimately be limited in precision at very nearly the same level that the nuclear result has currently reached.

Like neutron decay, pion beta decay has an advantage over nuclear decays in that there are no nuclear-structure-dependent corrections to be made. It also has the same advantage as the nuclear decays in being a purely vector transition, in its case $0^- \rightarrow 0^-$, so no separation of vector and axial-vector components is required. Its major disadvantage, however, is that the beta-decay mode, $\pi^+ \rightarrow \pi^0 e^+ \nu_e$, is a $\sim 10^{-8}$ branch of the total pion decay. This results in severe experimental limitations. For the pion beta decay, the value of V_{ud} is determined from the expression

$$V_{ud}^2 = \frac{K/\ln 2}{2G_F^2(1 + \Delta_R^V)f(1 + \delta_R)\tau_\pi/BR}, \quad (5)$$

where τ_π and BR are the mean life and branching ratio for pion beta decay, f is the statistical rate function and δ_R is the transition-dependent radiative correction evaluated for the case of the pion. The current world average [12] for the mean life is $\tau_\pi = 2.6033(5) \times 10^{-8}$ s. As for the beta-decay branching ratio, the world average is now dominated by the recent PIBETA measurement [16], which determined it relative to the much stronger $\pi^+ \rightarrow e^+ \nu_e(\gamma)$ branch and improved its experimental precision by more than a factor of 6. The latter branch has been determined experimentally but can also be determined – more precisely, in fact – by theoretical calculation [12].

With the theoretical value $BR(\pi^+ \rightarrow e^+ \nu_e(\gamma)) = 1.2352(5) \times 10^{-4}$, the PIBETA measurement of the pion beta-decay branching ratio becomes $1.040(6) \times 10^{-8}$. Equation 5 then yields the result

$$V_{ud} = 0.9751(27), \quad [\text{pion}] \quad (6)$$

which agrees well with the nuclear result but with 10 times the uncertainty. If, instead, the experimental measurement had been used for the $\pi^+ \rightarrow e^+ \nu_e(\gamma)$ branching ratio, then $V_{ud} = 0.9732(32)$, which also is consistent with the nuclear result but with an even larger uncertainty.

IV. CONCLUSIONS

As of now, superallowed $0^+ \rightarrow 0^+$ nuclear β decay clearly dominates in the determination of V_{ud} . A weighted average of the results quoted in Eqs. 2, 4 and 6, yields a result that differs only in the fifth place of decimals from the nuclear result alone; and, considering the ambiguities present in the other decays, especially in that of the neutron, it seems best for the time being to rely simply upon the unaveraged nuclear result. This result is not limited by experimental uncertainties but by uncertainties originating in the theoretical corrections applied to the data. Even so, the largest theoretical uncertainty is not due to the nuclear-structure-dependent corrections

but rather to the “inner” radiative correction, Δ_R^V , which actually is common to all three β decays: nuclear, neutron and pion. Clearly it would be beneficial to have confirmation of the nuclear V_{ud} by results of comparable precision from the other two β -decay modes, but one should not anticipate that either of the latter will actually surpass the nuclear result in the near future.

Although experiment is not currently limiting the precision of the nuclear V_{ud} , there is considerable activity at the moment in improving experimental precision and extending measurements to previously unstudied superallowed transitions. The approach is best illustrated by Fig. 1 and the observation that the calculated corrections are validated by their success in removing the ft -value scatter. Improvements in experimental precision would test the calculations even more severely, as would new examples of $0^+ \rightarrow 0^+$ transitions specifically selected for having *larger* calculated corrections. The reasoning is that if the ft values measured for cases with large calculated corrections also turn into corrected $\mathcal{F}t$ values that are consistent with the others, then this must verify the calculations’ reliability for the existing cases, which have smaller corrections. In fact, the cases of ^{34}Ar , ^{62}Ga and ^{74}Rb were chosen for this reason and their consistency has already led to considerably increased confidence in the calculated corrections.

It is reasonable to expect that there will be some further improvement in the nuclear value for V_{ud} as a result of this campaign to test the nuclear-structure-dependent corrections. However, really significant improvements must await a more precise calculation of Δ_R^V .

Acknowledgments

Much of the work reported here was done in collaboration with I.S. Towner. It was supported financially by the U.S. Department of Energy under Grant No. DE-FG03-93ER40773 and by the Robert A. Welch Foundation under Grant No. A-1397.

-
- [1] J.C. Hardy and I.S. Towner, Phys. Rev. C **71**, 055501 (2005); Phys. Rev. Lett. **94**, 092502 (2005).
 - [2] G. Savard *et al.*, Phys. Rev. Lett. **95**, 102501 (2005).
 - [3] I.S. Towner and J.C. Hardy, Phys. Rev. C **72**, 055501 (2005).
 - [4] T. Eronen *et al.*, Phys. Lett. B **636**, 191 (2006).
 - [5] T. Eronen *et al.*, Phys. Rev. Lett. **97**, 232501 (2006).
 - [6] G. Bollen *et al.*, Phys. Rev. Lett. **96**, 152501 (2006).
 - [7] P.H. Barker and A.P. Byrne, Phys. Rev. C **73**, 064306 (2006).
 - [8] V.E. Jacob *et al.*, Phys. Rev. C **74**, 055502 (2006).
 - [9] B. Hyland *et al.*, Phys. Rev. Lett. **97**, 102501 (2006).
 - [10] I.S. Towner and J.C. Hardy, Phys. Rev. C **66**, 035501 (2002).
 - [11] W.J. Marciano and A. Sirlin, Phys. Rev. Lett. **96**, 032002 (2006).
 - [12] W.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
 - [13] I.S. Towner and J.C. Hardy, J. Phys. G: Nucl. Part. Phys. **29** (2003) 197.
 - [14] J.S. Nico *et al.*, Phys. Rev. C **71**, 055502 (2005).
 - [15] A. Serebrov *et al.*, Phys. Lett. B **605**, 72 (2005).
 - [16] D. Počanić *et al.*, Phys. Rev. Lett. **93**, 181803 (2004).